



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search PubMed ☒ for ☒ Limits Preview/Index History Clipboard Details

About Entrez

 Abstract ☒ Show: 20 Text ☒

Text Version

☐ 1: J Protein Chem. 1999 Aug;18(6):619-25.

Related Articles, L

Entrez PubMed

Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services

Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources

Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

Oxidative folding of reduced and denatured huwentoxin-I.

Liang S, Shu Q, Wang X, Zong X.


College of Life Sciences, Hunan Normal University, Changsha, China.

Huwentoxin-I, a neurotoxic peptide with 33 amino acid residues and three disulfide bond was used to investigate the pathway of reduction/denaturation and of oxidative folding in small proteins with multiple disulfide bonds. Titration of thiol groups, reversed-phase HPLC, 1D NMR spectroscopy, and biological activity assays were used to monitor the extent of reduction/ denaturation and renaturation of the toxin. The reduction and denaturation of huwentoxin-I resulted in a 100% loss of bioactivity as measured in a mophrenic nerve-diaphragm preparation. About 90% of full biological activity could be restored under optimized conditions of oxidative refolding of the reduced peptide. Sever reaction conditions employing air oxidation, oxidized and reduced glutathione (GSSG and GSH), and cystine/cysteine were investigated in order to find optimal conditions for renaturation of huwentoxin-I. The best renaturation yield was achieved in 0.1 mM GSSG and 1 mM GSH at pH 8.5 and 4 degrees C over 24 hr. High concentrations of glutathion and high temperatures reduced renaturation yields. Oxidative refolding of huwentoxin-I air requires about 6 days for maximal yields and is inhibited by EDTA.

PMID: 10609637 [PubMed - indexed for MEDLINE]

 Abstract ☒ Show: 20 Text ☒[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act](#) | [Disclaimer](#)

Jan 29 2004 15:



Entrez PubMed Nucleotide Protein Genome Structure PMC Taxonomy Books

Search for

Limits Preview/Index History Clipboard Details
 Show:

☐ 1: P83591. Hainantoxin-I (Hn...[gi:32363268]

BLink, L

LOCUS P83591 33 aa linear INV 15-SEP-2003

DEFINITION Hainantoxin-I (HnTx-I).

ACCESSION P83591

VERSION P83591 GI:32363268

DBSOURCE swissprot: locus TXN1_SELHA, accession P83591;
 class: standard.
 created: Sep 15, 2003.
 sequence updated: Sep 15, 2003.
 annotation updated: Sep 15, 2003.

KEYWORDS Toxin; Neurotoxin; Ionic channel inhibitor; Sodium channel inhibitor; Amidation.

SOURCE Ornithoctonus hainana

ORGANISM Ornithoctonus hainana

Eukaryota; Metazoa; Arthropoda; Chelicerata; Arachnida; Araneae;
 Mygalomorphae; Theraphosidae; Ornithoctonus.

REFERENCE 1 (residues 1 to 33)

AUTHORS Li, D.-L., Xiao, Y.-C. and Liang, S.-P.

TITLE Direct Submission

JOURNAL Submitted (~MAY-2003) to Swiss-Prot

REMARK SEQUENCE, FUNCTION, SUBUNIT, SUBCELLULAR LOCATION, TISSUE
 SPECIFICITY, MASS SPECTROMETRY, DISULFIDE BONDS, AMIDATION, AND
 STRUCTURE BY NMR.
 TISSUE=Venom

COMMENT [FUNCTION] Is a depressant toxin. Binds and blocks insect sodium
 channels without altering the activation or inactivation kinetics.
 [SUBUNIT] Monomer.
 [SUBCELLULAR LOCATION] Secreted.
 [TISSUE SPECIFICITY] Expressed by the venom gland.
 [MASS SPECTROMETRY] MW=3608.01; METHOD=MALDI.
 [SIMILARITY] Belongs to the huwentoxin-I family.

FEATURES Location/Qualifiers
 source 1..33
 /organism="Ornithoctonus hainana"
 /db_xref="taxon:209901"
Protein 1..33
 /product="Hainantoxin-I"
Bond bond(2,17)
 /bond_type="disulfide"
Bond bond(9,22)
 /bond_type="disulfide"
Bond bond(16,29)
 /bond_type="disulfide"
Site 33
 /site_type="amidation"

ORIGIN
 1 eckgfgkscv pgkneccsgy acnsrdkwck vll
 //

h cb hg e e e e fcg b e



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search for

☒ Limits Preview/Index History Clipboard Details

About Entrez

☒ Show: ☒ Sort ☒ ☒

Text Version

☐ 1: Biochemistry. 1992 Feb 18;31(6):1749-56.

[Related Articles, L](#)

Entrez PubMed

Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services

Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources

Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

Disulfide exchange folding of insulin-like growth factor I.

Hober S, Forsberg G, Palm G, Hartmanis M, Nilsson B.

Department of Biochemistry and Biotechnology, Royal Institute of Technology, Stockholm, Sweden.

The disulfide exchange folding properties of insulin-like growth factor I (IGF-I) have been analyzed in a redox buffer containing reduced (10 mM) and oxidized (1 mM) glutathione. Under these conditions, the 3 disulfide bridges of the 70 amino acid peptide were not quantitatively formed. Instead, five major forms of IGF-I were detected, and these components were concluded to be in equilibrium as their relative amounts were similar starting from either reduced, native, or a mismatched variant of IGF-I containing two no native disulfides. The different components in the mixtures were trapped by thiol alkylation using vinylpyridine and subsequently isolated by reverse-phase HPLC. The purified variants were further characterized using plasma desorption mass spectrometry and peptide mapping. Two of the five different forms were identified as native and mismatched IGF-I. One form was a variant with only one disulfide bond, and the other two major components had two disulfides formed. In a separate experiment, early refolding intermediates were trapped by pyridylethylation after only 90 s of refolding in the glutathione buffer, starting from reduced IGF-I. The intermediates were identical to the components observed at equilibrium, but at different relative concentrations. On the basis of the disulfide bond patterns of the different components in the equilibrium mixtures, we conclude that the disulfide between cysteines-47 and -52 in IGF-I is an unfavorable high-energy bond that may exist in the native molecule in a strained configuration.

PMID: 1737028 [PubMed - indexed for MEDLINE]

☒ Show: ☒ Sort ☒ ☒


[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)

Jan 29 2004 15:



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search PubMed for

☒ Limits Preview/Index History Clipboard Details

About Entrez

Abstract Text

Text Version

☐ 1: Protein Sci. 1999 Aug;8(8):1605-13.

Related Articles, L

Entrez PubMed

Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services

Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources

Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

The disulfide-coupled folding pathway of apamin as derived from diselenide-quenched analogs and intermediates.

Pegoraro S, Fiori S, Cramer J, Rudolph-Bohner S, Moroder L.

Max-Planck-Institut für Biochemie, Martinsried, Germany.

The sequence of apamin, an 18 residue bee venom toxin, encloses all the information required for the correct disulfide-coupled folding into the cystine-stabilized alpha-helical motif. Three apamin analogs, each containing a pair of selenocysteine residues replacing the related cysteines, were synthesized to mimic the three possible apamin isomers with two crossed, parallel, or consecutive disulfides, respectively. Refolding experiments clearly revealed that the redox potential of selenocysteine prevails over the sequence encoded structural information for proper folding of apamin. Thus, selenocysteine can be used as new device to generate productive and nonproductive folding intermediates of peptides and proteins. In fact, disulfides are selectively reduced in presence of the diselenide and the conformational features derived from these intermediates as well as from the three-dimensional (3D) structures of the selenocysteine-containing analogs with their nonnatural networks of diselenide/disulfide bridges allowed to gain further insight into the subtle driving forces for the correct folding of apamin that mainly derive from local conformational preferences.

PMID: 10452604 [PubMed - indexed for MEDLINE]

Abstract Text


[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)

Jan 29 2004 15:



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search for

☒ Limits Preview/Index History Clipboard Details

About Entrez

Show: Sort:

Text Version

Entrez PubMed

Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services

Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources

Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

☐ 1: Arch Biochem Biophys. 1998 Jun 1;354(1):1-8.

Related Articles, L

ELSEVIER
FULL-TEXT ARTICLE

Unfolding/folding studies on cobrotoxin from Taiwan cobra venom: pH and GSH/GSSG govern disulfide isomerization at the C-terminus.

Chang LS, Lin SR, Chang CC.

Department of Biochemistry, Kaohsiung Medical College, Taiwan, Republic of China.
lschang@mail.nsysu.edu.tw

Refolding of cobrotoxin was assessed by the exposure degree of its single Trp determined by an acrylamide quenching study. The change in the accessibility of Trp for acrylamide quantitatively reflected the formation of folded cobrotoxin, and the data were confirmed by HPLC and gel electrophoresis analyses. However, the site-specific information provided by quenching Trp fluorescence revealed that the ordered structure in the neighborhood of Trp was attained prior to the complete formation of the tertiary structure of cobrotoxin. HPLC analyses showed that, in addition to refolded cobrotoxin, two novel species (cobrotoxin I and cobrotoxin III) with isomerization of disulfide bonds at the C-terminus of the toxin molecule were produced along the folding reaction. The disulfide pairings in cobrotoxin and cobrotoxin III were Cys43-Cys55 and Cys54-Cys60 and Cys43-Cys60 and Cys54-Cys55, respectively. Among the three possible two-disulfide species at the C-terminus, the two disulfide linkages Cys43-Cys60 and Cys54-Cys55 of cobrotoxin III caused a marked decrease in lethality and resulted in a conformation which was notably different from that observed with the native toxin molecule as evidenced by CD spectra. The refolding reaction was accelerated by the addition of GSH/GSSG, and the resulting products were mostly folded cobrotoxin. However, if GSH/GSSG was not added into the initial folding materials, the yields of cobrotoxin II and cobrotoxin III greatly increased. The conversion of cobrotoxin to its isomers was to be irreversible and pH-dependent: the higher the pH, the faster the rate of conversion. However, this conversion could be partly inhibited by GSH/GSSG. Cobrotoxin II and cobrotoxin III were purified from Taiwan cobra venom as well, and their yields in comparison to that of cobrotoxin in venom were similar to that noted with the folded products in the presence of GSH/GSSG. Moreover, the rate of disulfide isomerization was expected to be slow in venom fluid in which the pH was approximately pH 6.2. Thus, the finding that cobrotoxin represents the predominant neurotoxin species in Taiwan cobra venom is probably associated with the synergistic effects of GSH/GSSG and pH.

PMID: 9633591 [PubMed - indexed for MEDLINE]



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search PubMed for

☒ Limits Preview/Index History Clipboard Details

About Entrez

Display Summary

Text Version

Items 41-60 of 249

Previous of 13 Next

Entrez PubMed

Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities

PubMed Services











Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources

Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

- ☐ **41:** Ruoppolo M, Freedman RB. Related Articles, Li
 Refolding by disulfide isomerization: the mixed disulfide between ribonuclease T1 and glutathione as a model refolding substrate.
 Biochemistry. 1995 Jul 25;34(29):9380-8.
 PMID: 7626608 [PubMed - indexed for MEDLINE]
- ☐ **42:** Kim JI, Takahashi M, Ogura A, Kohno T, Kudo Y, Sato K. Related Articles, Li
 Hydroxyl group of Tyr13 is essential for the activity of omega-conotoxin GVIA, a peptide toxin for N-type calcium channel.
 J Biol Chem. 1994 Sep 30;269(39):23876-8.
 PMID: 7929033 [PubMed - indexed for MEDLINE]
- ☐ **43:** Hill JM, Alewood PF, Craik DJ. Related Articles, Li
 Conotoxin TVIIA, a novel peptide from the venom of Conus tulipa 2. Three-dimension solution structure.
 Eur J Biochem. 2000 Aug;267(15):4649-57.
 PMID: 10903497 [PubMed - indexed for MEDLINE]
- ☐ **44:** Frech C, Schmid FX. Related Articles, Li
 Influence of protein conformation on disulfide bond formation in the oxidative folding of ribonuclease T1.
 J Mol Biol. 1995 Aug 4;251(1):135-49.
 PMID: 7643382 [PubMed - indexed for MEDLINE]
- ☐ **45:** Sasaki T, Feng ZP, Scott R, Grigoriev N, Syed NI, Fainzilber M, Sato K. Related Articles, Li
 Synthesis, bioactivity, and cloning of the L-type calcium channel blocker omega-conotoxin TxVII.
 Biochemistry. 1999 Sep 28;38(39):12876-84.
 PMID: 10504258 [PubMed - indexed for MEDLINE]
- ☐ **46:** Gandia L, Lara B, Imperial JS, Villarroja M, Albillos A, Maroto R, Garcia AG, Olivera BM. Related Articles, Li
 Analogies and differences between omega-conotoxins MVIIC and MVIID: binding sites and functions in bovine chromaffin cells.
 Pflugers Arch. 1997 Dec;435(1):55-64.
 PMID: 9359903 [PubMed - indexed for MEDLINE]
- ☐ **47:** Kim JI, Takahashi M, Ohtake A, Wakamiya A, Sato K. Related Articles, Li
 Tyr13 is essential for the activity of omega-conotoxin MVIIC and GVIA, specific N-type calcium channel blockers.
 Biochem Biophys Res Commun. 1995 Jan 17;206(2):449-54.
 PMID: 7826361 [PubMed - indexed for MEDLINE]

- ☐ **48:** Dadlez M, Kim PS. Related Articles, Li
 Rapid formation of the native 14-38 disulfide bond in the early stages of BPTI folding. Biochemistry. 1996 Dec 17;35(50):16153-64. PMID: 8973187 [PubMed - indexed for MEDLINE]
- ☐ **49:** Corzo G, Adachi-Akahane S, Nagao T, Kusui Y, Nakajima T. Related Articles, Li
 Novel peptides from assassin bugs (Hemiptera: Reduviidae): isolation, chemical and biological characterization. FEBS Lett. 2001 Jun 22;499(3):256-61. PMID: 11423127 [PubMed - indexed for MEDLINE]
- ☐ **50:** Chung D, Gaur S, Bell JR, Ramachandran J, Nadasdi L. Related Articles, Li
 Determination of disulfide bridge pattern in omega-conopeptides. Int J Pept Protein Res. 1995 Sep-Oct;46(3-4):320-5. PMID: 8537186 [PubMed - indexed for MEDLINE]
- ☐ **51:** Maskos K, Huber-Wunderlich M, Glockshuber R. Related Articles, Li
 DsbA and DsbC-catalyzed oxidative folding of proteins with complex disulfide bridge patterns in vitro and in vivo. J Mol Biol. 2003 Jan 17;325(3):495-513. PMID: 12498799 [PubMed - indexed for MEDLINE]
- ☐ **52:** Guddat LW, Martin JA, Shan L, Edmundson AB, Gray WR. Related Articles, Li
 Three-dimensional structure of the alpha-conotoxin GI at 1.2 A resolution. Biochemistry. 1996 Sep 3;35(35):11329-35. PMID: 8784187 [PubMed - indexed for MEDLINE]
- ☐ **53:** Gray WR. Related Articles, Li
 Disulfide structures of highly bridged peptides: a new strategy for analysis. Protein Sci. 1993 Oct;2(10):1732-48. PMID: 8251945 [PubMed - indexed for MEDLINE]
- ☐ **54:** Mitchell SS, Shon KJ, Foster MP, Davis DR, Olivera BM, Ireland CM. Related Articles, Li
 Three-dimensional solution structure of conotoxin psi-PIIIE, an acetylcholine gated ion channel antagonist. Biochemistry. 1998 Feb 3;37(5):1215-20. PMID: 9477946 [PubMed - indexed for MEDLINE]
- ☐ **55:** Yamashita H, Hirose M. Related Articles, Li
 Oxidative regeneration and selective reduction of native disulfide bonds in the N-terminal half-molecule of ovotransferrin. J Biol Chem. 1993 Sep 5;268(25):19062-9. PMID: 8395527 [PubMed - indexed for MEDLINE]
- ☐ **56:** Sato K, Raymond C, Martin-Moutot N, Sasaki T, Omori A, Ohtake A, Kim JJ, Kohno T, Takahashi M, Seagar M. Related Articles, Li
 Binding of chimeric analogs of omega-conotoxin MVIIA and MVIIC to the N- and P/Q type calcium channels. FEBS Lett. 1997 Sep 8;414(2):480-4. PMID: 9315745 [PubMed - indexed for MEDLINE]
- ☐ **57:** Doughty SW, Blaney FE, Orlek BS, Richards WG. Related Articles, Li
 A molecular mechanism for toxin block in N-type calcium channels. Protein Eng. 1998 Feb;11(2):95-9.

PMID: 9605543 [PubMed - indexed for MEDLINE]

☐ **58:** [Pegoraro S, Fiori S, Cramer J, Rudolph-Bohner S, Moroder L.](#)

[Related Articles](#), [Li](#)



The disulfide-coupled folding pathway of apamin as derived from diselenide-quenched analogs and intermediates.

Protein Sci. 1999 Aug;8(8):1605-13.

PMID: 10452604 [PubMed - indexed for MEDLINE]

☐ **59:** [Chang LS, Lin SR, Chang CC.](#)

[Related Articles](#), [Li](#)



Unfolding/folding studies on cobrotoxin from Taiwan cobra venom: pH and GSH/GSS govern disulfide isomerization at the C-terminus.

Arch Biochem Biophys. 1998 Jun 1;354(1):1-8.

PMID: 9633591 [PubMed - indexed for MEDLINE]

☐ **60:** [Skalicky JJ, Metzler WJ, Ciesla DJ, Galdes A, Pardi A.](#)

[Related Articles](#), [Li](#)



Solution structure of the calcium channel antagonist omega-conotoxin GVIA.

Protein Sci. 1993 Oct;2(10):1591-603.

PMID: 8251934 [PubMed - indexed for MEDLINE]

Display Show: Sort

Items 41-60 of 249

Previous of 13 [Ne](#)

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)

Jan 29 2004 15:



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search for

☒ Limits Preview/Index History Clipboard Details

About Entrez

☒ Show: ☒

Text Version

☐ 1: J Pept Res. 2003 Apr;61(4):202-12.

[Related Articles, L](#)

Entrez PubMed
Overview
Help | FAQ
Tutorial
New/Noteworthy
E-Utilities



Detergent-assisted oxidative folding of delta-conotoxins.

DeLa Cruz R, Whitby FG, Buczek O, Bulaj G.

Department of Biology, University of Utah, Salt Lake City, Utah 84112, USA.

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Privacy Policy

Conotoxins comprise a diverse group of disulfide-rich peptides found in venoms of predatory Conus species. The native conformation of these peptides is marginally stable compared with alternative conformations, often resulting in low folding yields. The oxidative folding of hydrophobic delta-conotoxins was found to produce less than 1% of the native peptide [Bulaj, G. et al. (2001) Biochemistry 40, 13201]. In order to identify factors that might improve folding yields, we screened a number of additives including water-soluble polymers, detergents and osmolytes for their ability to increase steady-state accumulation of the native delta-conotoxin PVIA. The presence of a non-ionic detergent Tween and low temperature appeared to be the most effective factors in improving the oxidative folding. The detergent was also effective in promoting folding of other hydrophobic delta-conotoxins. Based on our findings, we discuss a possible mechanism for detergent-assisted folding and the general applicability of this mechanism to facilitating proper folding of hydrophobic, cysteine-rich peptides.

PMID: 12605605 [PubMed - indexed for MEDLINE]

☒ Show: ☒

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
[Freedom of Information Act](#) | [Disclaimer](#)

Jan 29 2004 15:



Entrez PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Book

Search for

☒ Limits Preview/Index History Clipboard Details

About Entrez

Show: Sort

Text Version

☐ 1: Biopolymers. 1996 Jun;38(6):733-44.

[Related Articles, L](#)

Entrez PubMed

[Overview](#)
[Help | FAQ](#)
[Tutorial](#)
[New/Noteworthy](#)
[E-Utilities](#)

PubMed Services

[Journals Database](#)
[MeSH Database](#)
[Single Citation Matcher](#)
[Batch Citation Matcher](#)
[Clinical Queries](#)
[LinkOut](#)
[Cubby](#)

Related Resources

[Order Documents](#)
[NLM Gateway](#)
[TOXNET](#)
[Consumer Health](#)
[Clinical Alerts](#)
[ClinicalTrials.gov](#)
[PubMed Central](#)

[Privacy Policy](#)

Oxidative folding of omega-conotoxin MVIIC: effects of temperature and salt.

Kubo S, Chino N, Kimura T, Sakakibara S.

Peptide Institute, Inc., Protein Research Foundation, Osaka, Japan.

Oxidative folding of omega-conotoxin MVIIC, a highly basic 26-amino acid peptide with three disulfide bonds, predominantly gave two products with mismatched disulfide bond in 0.1M NH₄OAc buffer (pH 7.7) at 21 degrees C both in the presence and absence of redox reagents such as reduced and oxidized glutathione. A low reaction temperature (5 degrees C) and a high salt concentration in buffer such as 2M (NH₄)₂SO₄ were necessary to obtain the correctly folded biologically active product. The folding reaction was found to proceed via a two-stage pathway of (I) the formation and (II) the rearrangement of the mismatched disulfide bonds. Both the reaction temperature and the salt strongly affected the equilibrium between mismatched and correctly formed disulfide bonds in the second stage. Such an effect of salts on the rearrangement reaction could be explained by anion binding at a low concentration and the salting out effect at a high concentration by analyzing the rank order of their effectiveness. The anion-binding effect was also confirmed by examining the folding of the tetra-acetylated peptide at the Lys side chains. CD study suggested that the yield of the biologically active product was correlated with conformational change as functions of temperature and salt concentration.

PMID: 8652794 [PubMed - indexed for MEDLINE]

Show: Sort

[Write to the Help Desk](#)

[NCBI | NLM | NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act | Disclaimer](#)

Jan 29 2004 15: